

Chiropractic and Mental Health: History and Review of Putative Neurobiological Mechanisms

Christopher Kent*

Director of Evidence-Informed Curriculum and Practice, Sherman College of Chiropractic, Spartanburg, South Carolina, USA

***Corresponding author:** Christopher Kent, Director of Evidence-Informed Curriculum and Practice, Sherman College of Chiropractic, Spartanburg, South Carolina, USA. Tel: +18645788770; Email: ckent@sherman.edu

Citation: Kent C (2018) Chiropractic and Mental Health: History and Review of Putative Neurobiological Mechanisms. Jou Neuro Psy An Brain Res: JNPB-103.

Received Date: 20 July, 2018; **Accepted Date:** 1 August, 2018; **Published Date:** 13 August, 2018

Abstract:

The chiropractic profession has a long history of acknowledging the relationship between nervous system function and mental health. This paper reviews the history of chiropractic involvement in mental health issues, chiropractic institutions specializing in the care of mental health problems, and the putative neurobiological mechanisms associated with vertebral subluxation and dysregulation of the autonomic nervous system.

1. Keywords

Chiropractic, history, mental health, vertebral subluxation, manipulation, depression, anxiety, addiction, hospitals, autonomic nervous system, biological oscillators, neuroplasticity, polyvagal theory, neurovisceral integration, heart rate variability, resiliency, adaptability, salutogenesis

2. Introduction

Musculoskeletal conditions are the predominant reason persons seek chiropractic care. The top five reported reasons for attending chiropractic care are low back pain/back pain, neck pain, extremity problems, wellness/maintenance and hip pain. The

top five reasons for pediatric cases to attend chiropractic care are musculoskeletal conditions, excessive crying, neurological conditions, gastrointestinal conditions, and ear, nose, and throat conditions [1]. Although many chiropractors and those they serve tend to focus on disorders associated with the physical body, abnormal nervous system function may also affect emotional and psychological health. The author completed a brief historical overview of chiropractic and mental health [2]. This work represents expansion of that paper, and inclusion of putative neurobiological mechanisms.

3. History

D.D. Palmer founded the chiropractic profession 123 years ago. He described vertebral subluxations as “slightly displaced vertebrae which press against nerves causing impingements, the result being too much or not enough functioning” [3]. According to his son, B.J. Palmer, “D.D. Palmer was the first man to discover that insanity was caused by displaced cervical vertebrae, that by replacing them the patient could be restored to normal condition” [4]. B.J. also described his expert testimony in a case where he stated, “If an atlas is subluxated it makes abnormal the functions of the brain.” In answer to the question, “What is to be done in insanity?” he admonished his reader to “Go back to cause. Adjust that and return that brain to its normal capacity and capability” [5]. Another pioneer in the field of mental health and chiropractic was attorney and chiropractor Willard Carver. Carver authored the book, *Psycho-Bio-Physiology*, and wrote, “Between the Psychology and the Physiology I have built the Biologic bridge that scientifically connects these two very important departments of human experience” [6].

In the 1920s, several inpatient mental health facilities were established where chiropractic adjustments were the dominant clinical service provided. Two of these were located in Davenport, Iowa. In 1922, the Chiropractic Psychopathic Sanitarium was established. The facility was later known as Forest Park Sanitarium. North Dakota Judge A. W. Ponath noted that at the North Dakota state mental hospital, the “cure and discharge rate” ranged from 18-27%, compared to 65% at Forest Park [7]. The second facility, Clear View Sanitarium, was established in 1926. In 1951, Clear View was acquired by the Palmer School of Chiropractic. Chiropractor W. Heath Quigley, who directed the sanitarium, described the clinical protocol: “Each day, each patient was examined with the neurocalometer (NCM). If the clinician interpreted the NCM to indicate nerve impingement, the patient was adjusted.” Quigley reported that the rooms were “sunny and bright,” and that meals included “large servings of fresh vegetables...from a garden” [8]. Unfortunately, both institutions closed, (Forest Park in 1959 and Clear View in 1961) in large measure because of third party pay issues. Insurance companies often refused to pay the costs of care.

Furthermore, Iowa statutes at the time did not provide for licensing specialized hospitals; only full service medical hospitals were eligible for licensure. Clear View was not licensed as a hospital, and functioned legally as a nursing home [9].

The 1970s saw a renewed interest in chiropractic care and mental health issues. In 1973, Chiropractor Herman S. Schwartz edited a book titled “Mental Health and Chiropractic: A Multidisciplinary Approach.” Contributors included Nobel Laureates Rene Dubos and Linus Pauling, and such notables as Scott Haldeman, A.E. Homewood, Joseph Janse, Alexander Lowen, and Thomas Szasz [10]. In 1949, Schwartz had published a preliminary report of 350 patients afflicted with a “nervous or mental disorder” and reported that the majority of them showed improvement under chiropractic care [11]. Schwartz was active in the ACA Council on Mental Health (formerly Council on Psychotherapy), which survived through the '70s, but no longer exists. In 1983, Quigley authored an article describing a four decades period where “treatment of the mentally ill was a highly motivated discipline within the chiropractic profession” [12]. In 1988, Goff wrote a review of the theory and practice of “chiropractic treatment for mental illness” [13]. Interest in this field continues. Blanks, Schuster and Dobson [14] published the results of a retrospective assessment of subluxation based chiropractic care on self related health, wellness and quality of life. This is, to the authors’ knowledge, the largest study of its kind ever undertaken regarding a chiropractic population. After surveying 2,818 respondents in 156 practices, a strong connection was found between persons receiving Network Spinal care (a chiropractic technique) and self reported improvement in health, wellness and quality of life.

A systematic review was published which examined psychological outcomes in randomized controlled trials of spinal manipulation. The study concluded that “There was some evidence that spinal manipulation improved psychological outcomes compared with verbal interventions...The clinical implications are that physical treatments, such as spinal manipulation have psychological benefits” [15]. Genthner et al [16] reported on a series of 15 patients with a history of depression. The Beck Depression Inventory II (BDI-II) was used to

measure the baseline level of depression and any post-chiropractic care changes following orthospinology care, a chiropractic technique focused on correcting misalignments of the craniocervical junction. A paired t-test demonstrated significant improvement in depression test scores. A study evaluating the role of chiropractic care in persons undergoing inpatient addiction care consisted of a three arm randomized clinical trial with two control groups (one receiving usual medical care, and the other placebo controlled). This was a single blind study utilizing subluxation-centered chiropractic care, Torque-Release technique, implemented in a residential addiction care setting. The active group showed a significant decrease in anxiety while the placebo group showed no decrease in anxiety [17]. Other articles addressing mental health issues and chiropractic care have been published, ranging from single case reports to randomized clinical trials. Favorable responses were reported in persons with conditions including depression [18], ADHD [19], autism [20], dyslexia and learning disabilities [21]. Additionally, published papers report changes in general health measures in chiropractic patients using the RAND-36 and Global Well Being Scale (GWBS) [22], changes in domains of health related quality of life among public safety personnel undergoing chiropractic care [23], and chiropractic care in patients with cancer-related traumatic stress symptoms [24].

4. Salutogenesis

Chiropractic care incorporates a salutogenic approach. Sociologist Aaron Antonovsky coined the term salutogenesis in 1979. It is derived from *salus*, Latin for health, and *genesis*, meaning to give birth. Salutogenesis, the study of the origins and creation of health, provides a method to identify an interconnected way to enhance well-being. Salutogenesis provides a framework for a method of practice to promote health [25].

Salutogenic theory goes to the very essence of neurobiology. It has been noted that neurological processes (as well as anatomical structures) are remodeled by sensory input. These processes, collectively termed neuroplasticity, are operative at all levels of the nervous system. Smith [26] described the range of these mechanisms: "From the afferent (incoming) activity of peripheral sensory receptors to

the efferent (outgoing) activity directed toward neuroendocrine organs, blood vessels, and muscles. Although the selectivity of perception probably makes it impossible to be aware of everything that is happening throughout the body, it is evident that these regulatory processes are essential for one's health, and that they provide the basis for functional salutogenic mechanisms of the brain." Smith further noted, "An organism with a salutogenic brain would experience the world as manageable and coherent ... with a self-perpetuating cycle for enhancing self-confidence and well-being."

5. Stress Responsivity

Hans Selye [27] pioneered investigations of the biological effects of stress in 1936 with the publication of his paper, "A syndrome produced by diverse noxious agents." Since then, more than 100,000 articles and books have been written on the subject. Selye describes stress as the nonspecific response to any demand. Although many individuals have concluded that stress is inevitably destructive, this view is incorrect. Selye noted, "Stress is not necessarily bad for you. It is also the spice of life, for any emotion, any activity causes stress...the same stress that makes one person sick is an invigorating experience for another...Complete absence of stress is incompatible with life since only a dead man makes no demand on his body or mind." Selye described two types of stress: Dis-stress -- from the Latin "bad," as in dissonance, and Eu-stress from the Greek "true" or "good," as in eutonia. Whether we experience a pleasant or unpleasant result from an event depends upon how our nervous system perceives, processes, and interprets that event. More than 15 years before Selye's historic publication, B.J. Palmer and J.H. Craven [28] described a similar concept: concussion of forces. This term refers to the meeting of external invasive forces and internal resistive forces. Just as stress may be destructive or beneficial, concussion of forces may produce or reduce vertebral subluxation. The result is dis-ease or ease. "That which caused the normal cycle to become abnormal was a concussion of forces centering at some point in the spinal column causing a subluxation...tissues do not nor cannot express their normal function." Palmer [29] quotes Webster's definition of adaptation: "To make suitable; to fit; or suit; to adjust; alter so as to fit for a new use." More than 60 years later, Selye [30] wrote,

"Every living being has a certain innate amount of adaptation energy or vitality." When a concussion of forces is corrective, Palmer [29] noted the following changes: "Perversion changed to verification; abuse to proper natural use; abnormal interpretation to normal interpretation; distortion to healthful manifestation; corruption to correction." Although it is unlikely that Selye was familiar with the writings of Palmer and Craven, the similarities are striking: Stress and concussion of forces; eu-stress and ease; dis-stress and disease. The practical application of these concepts requires a working definition of health. The World Health Organization (WHO) [31] defines health as "A state of complete physical, mental, and social well-being and not merely the absence of disease or infirmity." In this context, Selye [30] wrote, "The secret of health and happiness lies in successful adjustment to the ever-changing conditions on this globe; the penalties for failure in this great process of adaptation are disease and unhappiness."

6. Putative Neurobiological Mechanisms

6.1. Vertebral Subluxation

In 1906, DD Palmer and BJ Palmer [32] defined subluxation as follows: "A (sub)luxation of a joint, to a chiropractor, means pressure on nerves, abnormal functions creating a lesion in some portion of the body, either in its action, or makeup". Lantz [33] noted, "Common to all concepts of subluxation are some form of kinesiologic dysfunction and some form of neurologic involvement". Mechanical and degenerative changes associated with vertebral subluxation may result in a variety of neurological consequences:

- **Cord compression and adverse cord tension:** Compression of the spinal cord may result from disc protrusion, ligamentum flavum hypertrophy/corrugation, or osteophytosis. Myelopathy may result in cord pressure and/or pressure which interferes with the arterial supply [34-39].
- **Nerve root compression:** Compromise of the nerve roots may develop following disc protrusion or osteophytosis [40]. Spinal

nerve roots are exquisitely sensitive to compression [41-43].

- **Local irritation:** This includes irritation of mechanoreceptive and nociceptive fibers within the intervertebral motion segments [44].
 - **Vertebral artery compromise:** MacNab advises that osteophytes may cause vertebral artery compression [45].
 - **Autonomic dysfunction:** Symptoms associated with the autonomic nervous system have been reported in patients with cervical spine trauma. The Barre'-Lieou syndrome includes blurred vision, tinnitus, vertigo, temporary deafness, and shoulder pain. This phenomenon is also known as the posterior cervical syndrome [46] Stimulation of sympathetic nerves has been implicated in the pathogenesis of this syndrome [47].
 - **Coherence and oscillatory patterns:** Coherent oscillations are a characteristic of the human brain. [48] Furthermore, it has been proposed that synchronization of multiple rhythms is an essential manifestation of living processes [49]. Epstein describes wave activity association with Network Spinal care, a chiropractic technique involving light touches to the spine. According to Senzon, Epstein and Lemberger, "The network wave occurs at a higher self-organizational threshold, in the absence of significant adverse mechanical cord tension, and with enhanced self-regulation of the spinal subsystems. With the onset of central pattern generation, modulated through the network wave, reorganizational behavior may emerge in the individual's spine and life as a whole" [50].
- ### 6.2. Operational Model of Vertebral Subluxation
- The author has proposed an operational model for the assessment of neurological dysregulation associated with vertebral subluxation [51]. The four components of this model include:
- **Dysafferentation:** The intervertebral motion segment is richly endowed with nociceptive and

mechanoreceptive structures [52-57]. As a consequence, biomechanical dysfunction caused by vertebral subluxation may result in altered nociception and/ or mechanoreception.

- **Dyskinesia:** Dyskinesia refers to distortion or impairment of voluntary movement [58]. Spinal motion may be reliably measured using inclinometry [59]. Alterations in regional ranges of motion may be associated with vertebral subluxation [60].
- **Dysponesis:** Dysponesis is evidenced by abnormal tonic muscle activity. Dysponesis refers to a reversible physiopathologic state consisting of errors in energy expenditure, which is capable of producing functional disorders. Dysponesis consists mainly of covert errors in action potential output from the motor and premotor areas of the cortex and the consequences of that output. These neurophysiological reactions may result from responses to environmental events, bodily sensations, and emotions. The resulting aberrant muscle activity may be evaluated using surface electrode techniques [61,62]. Typically, static surface electromyography (sEMG) with axial loading of the spine is used to evaluate innate responses to gravitational stress [63].
- **Dysautonomia:** The autonomic nervous system regulates the actions of organs, glands, and blood vessels. Acquired dysautonomia may be associated with a broad array of functional abnormalities [64-70]. Sympathetic tone may be evaluated by measuring skin temperature differentials using paraspinal infrared thermography [71]. Such techniques have been used to monitor changes in neurological function associated with vertebral subluxations [72].

7. Autonomic Dysregulation and Mental Health

Variability in heart rate reflects the vagal and sympathetic function of the autonomic nervous system, and is used as a monitoring tool in clinical conditions characterized by altered autonomic nervous system activity. Spectral analysis of beat-to-beat variability is a simple, non-invasive technique to evaluate autonomic dysfunction. Vertebral subluxations are changes in the position or motion of

a vertebra, which result in the interference with nerve function. Vertebral subluxations may result in altered autonomic nervous system activity. Heart rate variability is a reliable and valid tool that may be used to assess the changes in autonomic activity associated with the reduction and correction of vertebral subluxations [72]. Recent studies have reported the potential utility of HRV in the evaluation of conditions and states associated with autonomic dysregulation. These include carotid intima media thickness [73], prediction of mortality [74], multiple sclerosis [75,76], eating behavior [77], burnout and depression [78], chronic posttraumatic stress disorder [79], working memory performance [80], dementia [81], inflammation in rheumatoid arthritis [82], insulin resistance and metabolic syndrome [83], type 1 diabetes [84], cardiac autonomic nerve function in obese school-age children [85], cancer prognosis [86,87] and cognition [88,89]. In the mental health field, associations have been identified between cardiac vagal activity, immunometabolic risk factors, and depression [90]. Higher Beck Depression Inventory-II (BDI-II) scores were associated with decreased HRV [91]. Oh and Chae [92] note that HRV may be a crucial marker for mental health. They report that “HRV properties might be related to the degree of optimistic perspectives on life, and suggests that HRV markers of autonomic nervous system function could reflect positive human mind states.” Fiskum et al [93] state, “Internal psychopathology and dysregulated negative affect are characterized by dysregulation in the autonomic nervous system and reduced heart rate variability (HRV) due to increases in sympathetic activity alongside reduced vagal tone...Higher informational entropy was related to less psychopathology and less negative effect, and may provide an index of the organizational flexibility of the neurovisceral system.”

Polyvagal theory (PVT), proposed by Porges [94] posits that physiological state limits the range of behavior and psychological experience. Porges notes, “The theory links the evolution of the autonomic nervous system to affective experience, emotional expression, facial gestures, vocal communication, and contingent social behavior. In this way, the theory provides a plausible explanation for the reported covariation between atypical autonomic regulation (eg, reduced vagal and increased

sympathetic influences to the heart) and psychiatric and behavioral disorders that involve difficulties in regulating appropriate social, emotional, and communication behaviors.” Sullivan et al [95] explain that “PVT links the evolution of the autonomic nervous system to the emergence of prosocial behaviors and posits that the neural platforms supporting social behavior are involved in maintaining health, growth and restoration. This explanatory model which connects neurophysiological patterns of autonomic regulation and expression of emotional and social behavior, is increasingly utilized as a framework for understanding human behavior, stress and illness.” The authors describe how PVT is related to self-regulation, resilience, and adaptability. Smith et al [96] proposed the neurovisceral integration (NVI) model to explain observed relationships between peripheral physiology, cognitive performance, and emotional and physical health. This model is supported largely from studies examining cardiac vagal control. An expanded model describes the multilevel structure and function of vagal control. Higher levels are associated with cognitive/attentional responses, regulation based on perceptual representation of one's current somatic/visceral state, regulation based on conceptualization of sensory input and past experience, and amplifying, maintaining, or suppressing representations based on current goals. In reviewing the literature concerning HRV and chiropractic care, Kent concluded, “Case reports suggest that favorable changes in heart rate variability may follow reduction or correction of vertebral subluxations. Higher quality studies of larger populations should be conducted. It is biologically plausible that the changes in autonomic nervous system function following reduction or correction of vertebral subluxation may be objectively assessed using heart rate variability” [72].

8. Conclusion

Chiropractic care is concerned with the totality of the human experience. Vertebral subluxations may result in autonomic dysregulation, compromising the adaptive capacity of the organism. By analyzing and correcting vertebral subluxations, a patient is placed on a more optimum physiological path, potentially increasing resilience and adaptability. Further

research into the effects of vertebral subluxations on mental health, the neurobiological mechanisms involved, and the use of reliable and valid outcomes assessments should be undertaken. It is biologically plausible that vertebral subluxations compromise nervous system function and affect mental health.

References

1. Beliveau PJH, Wong JJ, Sutton DA, Simon NB, Bussièrès AE, et al. (2017) The chiropractic profession: a scoping review of utilization rates, reasons for seeking care, patient profiles, and care provided. *Chiropr Man Therap* 25: 35.
2. Kent C (2013) Chiropractic and mental health: a brief overview. *Journal of Philosophy, Principles & Practice of Chiropractic* 1-3.
3. Palmer DD (1910) *The Chiropractor's Adjuster: Text-book of the Science, Art and Philosophy of Chiropractic for Students and Practitioners.* Portland Oregon: Portland Printing House Company.
4. Palmer BJ (1905) *History Repeats. The Palmer School of Chiropractic.* Davenport, IA. 1951. Quoting from the *Chiropractor* 1.
5. Palmer BJ (1920) *The Science of Chiropractic. Volume 2. The Palmer School of Chiropractic.* Davenport, IA 41.
6. Carver W (1920) *Psycho-Bio-Physiology.* Book Department. Carver Chiropractic College. Oklahoma City.
7. Editorial in *The Times.* Westminster, MD. 1/31/36. P. 16. Quoted by Keating.
8. Quigley WH (1910) *Clear View Sanitarium Part 5. Dynamic Chiropractic* 8: 8.
9. Quigley WH (1992) *Clear View Sanitarium The final years. Dynamic Chiropractic* 10: 13.
10. Schwartz HS (ed) (1973) *Mental Health and Chiropractic: A Multidisciplinary Approach.* Sessions Publishers. New York.
11. Schwartz HS (1949) Preliminary analysis 350 mental patients' records treated by chiropractors. *Journal of National Chiropractic Association* 12-15.
12. Quigley WH (1983) Pioneering mental health: institutional psychiatric care in chiropractic. *Chiropractic History* 3: 69-73.
13. Goff P (1988) Chiropractic treatment of mental illness: a review of theory and practice. *Research Forum* 4: 4-10.
14. Blanks RHI, Schuster TL, Dobson M (1997) A retrospective assessment of Network care using a survey of self reported health, wellness and

- quality of life. *Journal of Vertebral Subluxation Research* 1:15.
15. Williams NH, Hendry M, Lewis R, Russell I, Westmoreland A, et al. (2007) Psychological response in spinal manipulation (PRISM): A systematic review of psychological outcomes in randomized controlled trials. *Complementary Therapies in Medicine* 15: 271-283.
 16. Genthner GC, Friedman HL, Studley CF (2005) Improvement in depression following reduction of upper cervical vertebral subluxation using orthospinology technique. *Journal of Vertebral Subluxation Research*.
 17. Holder JM, Duncan Robert C, Gissen M, Miller M, Blum K (2001) Increasing retention rates among the chemically dependent in residential treatment: auriculotherapy and (in a separate study) subluxation-based chiropractic care. *Journal of Molecular Psychiatry* 6.
 18. Desaulniers AMJ (2008) Effect of subluxation-based chiropractic care on quality of life in a patient with major depression. *Journal of Vertebral Subluxation Research*.
 19. Lovett L, Blum CL (2006) Behavioral and learning changes secondary to chiropractic care to reduce subluxations in a child with Attention Deficit Hyperactivity Disorder: A case study. *Journal of Vertebral Subluxation Research*.
 20. Khorshid KA, Sweat RW, Zemba DA, Zemba BN (2006) Clinical efficacy of upper cervical versus full spine chiropractic care on children with autism: A randomized clinical trial. *Journal of Vertebral Subluxation Research*.
 21. Pauli Y (2007) The effects of chiropractic care on individuals suffering from learning disabilities and dyslexia: A review of the literature. *Journal of Vertebral Subluxation Research*.
 22. Blanks RHI, Dobson M (1999) A study regarding measures of general health status in patients using the Bio Energetic Synchronization Technique: A follow up study. *Journal of Vertebral Subluxation Research* 3: 1.
 23. McAllister W, Boone WR (2007) Changes in physical state and self-perceptions in domains of health related quality of life among public safety personnel undergoing chiropractic care. *Journal of Vertebral Subluxation Research*.
 24. Monti DA, Stoner ME, Zivin G, Schlesinger M (2007) Short term correlates of the Neuro Emotional Technique for cancer-related traumatic stress symptoms: A pilot case series. *J Cancer Surviv* 1: 161-166.
 25. Becker CM, Glascoff MA, Felts WM, Kent C (2015) Adapting and using quality management methods to improve health promotion. *Explore (NY)* 11: 222-228.
 26. Smith DF (2002) Functional salutogenic mechanisms of the brain. *Perspectives in Biology and Medicine* 45: 319-328.
 27. Selye H (1936) A syndrome produced by diverse noxious agents. *Nature* 138: 32.
 28. Palmer BJ, Craven JH (1920) *The Philosophy of Chiropractic*. Davenport, IA. Palmer School of Chiropractic.
 29. Palmer BJ (1951) *History Repeats*. Davenport, IA. Palmer School of Chiropractic.
 30. *The Stress of Life*. New York. McGraw Hill, Co. 1984.
 31. *The first ten years of the World Health Organization*. World Health Organization. Geneva. 1958.
 32. Palmer DD, Palmer BJ (1906) *The Science of Chiropractic*. Davenport, IA: The Palmer School of Chiropractic.
 33. Lantz CA (1995) The subluxation complex. In: Gatterman MI, ed. *Foundations of Chiropractic Subluxation*. St. Louis, MO: Mosby.
 34. O'Connell JE (1955) Involvement of the spinal cord by intervertebral disc protrusions. *Br J Surg* 43: 225.
 35. Taylor AR (1953) Mechanism and treatment of spinal cord disorders associated with cervical spondylosis. *Lancet* 1: 717.
 36. Mair WG, Druckman R (1953) The pathology of spinal cord lesions and their relations to the clinical features in protrusion of cervical intervertebral discs. *Brain* 76:70-91.
 37. Maiuri F, Gangemi M, Gambardella A, Simari R, D'Andrea F (1985) Hypertrophy of the ligamenta flava of the cervical spine. Clinico-radiological correlations. *J Neurosurg Sci* 29: 89-92.
 38. Breig A (1970) Overstretching of and circumscribed pathological tension in the spinal cord--a basic cause of symptoms in cord disorders. *J Biomech* 3:7-9.
 39. Rydevik BL (1992) The effects of compression on the physiology of nerve roots. *J Manipulative Physiol Ther* 15: 62-66.
 40. MacNab I (1975) Cervical spondylosis. *Clin Orthop* 109: 69-77.
 41. Ando M, Tamaki T, Kawakami M, Minamide A, Nakagawa Y, et al (2013) Electrophysiological diagnosis using sensory nerve action potential for the intraforaminal and extraforaminal L5 nerve root entrapment. *Eur Spine J* 22: 833-839.

42. Kobayashi S, Yoshizawa H, Yamada S (2004) Pathology of lumbar nerve root compression. Parts 1 and 2. *Journal of Orthopedic Research* 22: 170-188.
43. Sharpless, SK (1975) Susceptibility of spinal roots to compression block. NINCDS Monograph 15, DHEW publication (NIH) 155-161.
44. Kent C (1996) Models of vertebral subluxation: a review. *Journal of Vertebral Subluxation Research*. August 1: 1-7.
45. MacNab I (1975) Cervical spondylosis. *Clin Orthop* 109: 69-77.
46. Barre' JA (1926) Sur un syndrome sympathique cervical posterieur et sa cause frequente, 1, artrite cervicale. *Rev Neurol (Paris)* 1: 1246-1248.
47. Watanuki A (1981) (The effect of the sympathetic nervous system on cervical spondylosis). *Nippon Seikeigeka Gakkai Zasshi* 55: 371-385.
48. Akam TE, Kullmann DM (2012) Efficient "communication through coherence" requires oscillations structured to minimize interference between signals. *PLoS Comput Biol* 8: e1002760.
49. Muehsam D, Ventura C (2014) Life rhythm as a symphony of oscillatory patterns: electromagnetic energy and sound vibration modulates gene expression for biological signaling and healing. *Glob Adv Health Med* 3: 40-55.
50. Senzon SA, Epstein DM, Lemberger D (2016) The Network spinal wave as a central pattern generator. *J Altern Complement Med* 22: 544-556.
51. Kent C (2011) A four-dimensional model of vertebral subluxation. *Dynamic Chiropractic*.
52. Bogduk N, Tynan W, Wilson AS (1981) The nerve supply to the human lumbar intervertebral discs. *J Anat* 132: 39-56.
53. Nakamura S, Takahashi K, Takahashi Y, Shimada Y, Moriya H (1996) Origin of nerves supplying the posterior portion of lumbar intervertebral discs. *Spine* 21: 917-924.
54. Mendel T, Wink CS, Zimny ML (1992) Neural elements in human cervical intervertebral discs. *Spine* 17: 132-135.
55. McLain RF (1994) Mechanoreceptor endings in human cervical facet joints. *Spine* 19: 495-501.
56. Jiang H, Russell G, Raso VJ, Moreau MJ, Hill DL, et al. (1995) The nature and distribution of the innervation of human supraspinal and interspinal ligaments. *Spine* 20: 869-876.
57. Rhalmi S, Yahia LH, Newman N, Isler M (1993) Immunohistochemical study of nerves in lumbar spine ligaments. *Spine* 18: 264-267.
58. Dorland's Pocket Medical Dictionary. 25th edition. WB Saunders Company. 1995.
59. Saur PM, Ensink FB, Frese K, Seeger D, Hildebrandt J (1996) Lumbar range of motion: reliability and validity of the inclinometer technique in the clinical measurement of trunk flexibility. *Spine* 21: 1332-1338.
60. Blunt KL, Gatterman MI, Bereznick DE (1995) Kinesiology: an essential approach toward understanding the chiropractic subluxation. Chapter 11. In: Gatterman MI (ed): *Foundations of Chiropractic Subluxation*. Mosby, St. Louis, MO.
61. Whatmore GB, Kohi DR (1968) Dysponesis: a neurophysiologic factor in functional disorders. *Behav Sci* 13: 102-124.
62. Large R, Butler M, James F, Peters J (1990) A systems model of chronic musculo-skeletal pain. *Aust N Z J Psychiatry* 24: 529-536.
63. Kent C (1997) Surface electromyography in the assessment of changes in paraspinal muscle activity associated with vertebral subluxation: a review. *Journal of Vertebral Subluxation Research* 1:15.
64. Backonja M-M (1994) Reflex sympathetic dystrophy/sympathetically mediated pain/causalgia: the syndrome of neuropathic pain with dysautonomia. *Seminars in Neurology* 14: 263.
65. Goldstein DS, Holmes C, Cannon III RO, Eisenhofer G, Kopin IJ (1997) Sympathetic cardioneuropathy in dysautonomias. *New Engl J Med* 336: 696-702.
66. Vassallo M, Camilleri M, Caron BL, Low PA (1991) Gastrointestinal motor dysfunction in acquired selective cholinergic dysautonomia associated with infectious mononucleosis. *Gastroenterology* 100: 252-258.
67. Baron R, Engler F (1996) Postganglionic cholinergic dysautonomia with incomplete recovery: a clinical, neurophysiological and immunological case study. *J Neurol* 243: 18.
68. Soares JLD (1996) Dysautonomias. *Acta Medica Portuguesa* 8: 425.
69. Stryes KS (1994) The phenomenon of dysautonomia and mitral valve prolapse. *J Am Acad Nurse Practitioners* 6: 11.
70. Uematsu S, Edwin DH, Jankel ER, Kozikowski J, Trattner M (1988) Quantification of thermal asymmetry. *J Neurosurg* 69: 552.

71. Miller JL (1964) Skin temperature differential analysis. *International Review of Chiropractic (Science)* 1: 41.
72. Kent C (2017) Heart rate variability to assess the changes in autonomic nervous system function associated with vertebral subluxation. *Res Rev Neurosci* 1: 14-21.
73. Pereira VL Jr, Dobre M, Dos Santos SG, Fuzatti JS, Oliveira CR, et al. (2017) Association between carotid intima media thickness and heart rate variability in adults at Increased cardiovascular risk. *Front Physiol* 8: 248.
74. Lee CH, Lee JH, Son JW, Kim U, Park JS, et al. (2018) Normative values of short-term heart rate variability parameters in Koreans and their clinical value for the prediction of mortality. *Heart Lung Circ* 27: 576-587.
75. Vleck M, Penesova A, Imrich R, Meskova M, Mravcova M, et al. (2017) Autonomic nervous system response to stressors in newly diagnosed patients with multiple sclerosis. *Cell Mol Neurobiol* 38: 363-370.
76. Studer V, Rocchi C, Motta C, Lauretti B, Perugini J, et al. (2017) Heart rate variability is differentially altered in multiple sclerosis: implications for acute, worsening and progressive disability. *Mult Scler J Exp Transl Clin* 3:2055217317701317.
77. Ozpelit ME, Ozpelit E (2017) How we eat may be as important as what we eat: eating behaviour and heart rate variability. *Acta Cardiol* 72: 299-304.
78. Kanthak MK, Stalder T, Hill, Thayer JF, Penz M et al. (2017) Autonomic dysregulation in burnout and depression: evidence for the central role of exhaustion. *Scand J Work Environ Health* 43: 475-484.
79. Park JE, Lee JY, Kang SH, Choi JH1, Kim TY, et al. (2017) Heart rate variability of chronic posttraumatic stress disorder in the Korean veterans. *Psychiatry Res* 255:72-77.
80. Giuliano RJ, Gatzke-Kopp LM, Roos LE, Skowron EA (2017) Resting sympathetic arousal moderates the association between parasympathetic reactivity and working memory performance in adults reporting high levels of life stress. *Psychophysiology* 54: 1195-1208.
81. da Silva VP, Oliveira BRR, Mello RGT, Moraes H, Deslandes AC, et al. (2018) Heart rate variability indexes in dementia: a systematic review with a quantitative analysis. *Curr Alzheimer Res* 15: 80-88.
82. Koopman FA, van Maanen MA, Vervoordeldonk MJ, Tak PP (2017) Balancing the autonomic nervous system to reduce inflammation in rheumatoid arthritis. *J Intern Med* 282: 64-75.
83. Saito I, Maruyama K, Eguchi E, Kato T, Kawamura R, et al. (2017) Low heart rate variability and sympathetic dominance modifies the association between insulin resistance and metabolic syndrome - The Toon Health Study. *Circ J* 81: 1447-1453.
84. Silva AKFD, Christofaro DGD, Bernardo AFB, Vanderlei FM, Vanderlei LCM, et al. (2017) Sensitivity, specificity and predictive value of heart rate variability indices in type 1 diabetes mellitus. *Arq Bras Cardiol* 108: 255-262.
85. Yi LF, Wen HX, Huang XL, Qiu M, Cao XX (2017) [Cardiac autonomic nerve function in obese school-age children.] Article in Chinese. *Zhongguo Dang Dai Er Ke Za Zhi* 19: 524-528.
86. Wang YM, Wu HT, Huang EY, Kou YR, Hseu SS (2013) Heart rate variability is associated with survival in patients with brain metastasis: a preliminary report. *Biomed Res Int* 2013: 503421.
87. Guo Y, Koshy S, Hui D, Palmer JL, Shin K, et al. (2015) Prognostic value of heart rate variability in patients with cancer. *J Clin Neurophysiol* 32: 516-520.
88. Zeki Al Hazzouri A, Haan MN, Deng Y, Neuhaus J, Yaffe K (2014) Reduced heart rate variability is associated with worse cognitive performance in elderly Mexican Americans. *Hypertension* 63: 181-187.
89. Frewen J, Finucane C, Savva GM, Boyle G, Coen RF et al. (2013) Cognitive function is associated with impaired heart rate variability in ageing adults: the Irish longitudinal study on ageing wave one results. *Clin Auton Res* 23: 313-323.
90. Hu MX, Penninx BWJH, de Geus EJC, Lamers F, Kuan DC, et al. (2018) Associations of immunometabolic risk factors with symptoms of depression and anxiety: the role of cardiac vagal activity. *Brain Behav Immun*.
91. Huang M, Shah A, Su S, Goldberg J, Lampert RJ, et al. (2018) Association of depressive symptoms and heart rate variability in Vietnam war-era twins: a longitudinal twin difference Study. *JAMA Psychiatry* 75: 705-712.
92. Oh J, Chae JH (2018) Linear and nonlinear dynamics of heart rate variability are correlated with purpose in life and degree of optimism in anxiety disorder patients. *Nonlinear Dynamics Psychol Life Sci* 22: 173-190.

93. Fiskum C, Andersen TG, Bornas X, Aslaksen PM, Flaten MA, et al. (2018) Non-linear heart rate variability as a discriminator of internalizing psychopathology and negative affect in children with internalizing problems and healthy controls. *Front Physiol* 9: 561.
94. Porges SW (2009) The polyvagal theory: New insights into adaptive reactions of the autonomic nervous system. *Cleve Clin J Med* 76: S86–S90.
95. Sullivan MB, Erb M, Schmalzl L, Moonaz S, Noggle Taylor J, et al. (2018) Yoga therapy and polyvagal theory: the convergence of traditional wisdom and contemporary neuroscience for self-regulation and resilience. *Front Hum Neurosci* 12: 67.
96. Smith R, Thayer JF, Khalsa SS, Lane RD (2017) The hierarchical basis of neurovisceral integration. *Neurosci Biobehav Rev* 75: 274-296.

Copyright: ©2018 Christopher Kent *. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permit unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.